

Medication	EMT	AEMT	I	P	Route
Acetaminophen	•	•	•	•	PO
Adenosine			•	•	IV/IO
Afrin	•	•	•	•	IN
Aspirin	•	•	•	•	PO
Albuterol Sulfate	•	•	•	•	Inhaled
Atropine Sulfate			•	•	IV/IO
Calcium Chloride			•	•	IV/IO
Dextrose 10% / 50%		•	•	•	IV/IO
Diphenhydramine (Benadryl)	•	•	•	•	PO
Diphenhydramine (Benadryl)		•	•	•	IM/IV
Dobutamine			•	•	IV/IO
Enalapril			•	•	IV/IO
Epinephrine 1:1000	•	•	•	•	IM <b>Epi Pen ONLY</b>
Epinephrine 1:1000			•	•	Calculated IM
Epinephrine 1:10,000			•	•	IV/IO
Glucagon		•	•	•	IM/IN
Glucagon			•	•	Calculated IV <sub>1</sub>
Ketorolac (Toradol)			•	•	IM/IV/IO
Lidocaine			•	•	IV/IO

Magnesium Sulfate 50%			•	•	IV/IO
Methylprednisone			•	•	IM/IV/IO
Midazolam (Versed)			•	•	IM/IV/IO/IN
Morphine Sulfate			•	•	IM/IV/IO
Naloxone (Narcan)	•				<b>IN only with a fixed/unit dosed medication</b>
Naloxone (Narcan)		•	•	•	<b>IV/IO/IN Dose calculation/measurement</b>
Nitroglycerin SL	•				<b>SL with permission from Medical Control</b>
Nitroglycerin SL		•	•	•	SL
Nitroglycerin Paste		•	•	•	Transdermal
Norepinephrine (Levophed)			•	•	IV/IO Infusion
Ondansetron (Zofran) ODT	•	•	•	•	PO (Orally Disolvable Tablet)
Ondansetron (Zofran) Vial		•	•	•	IM/IV/IO
Predisolone			•	•	PO
Sodium Bicarbonate			•	•	IV/IO
*** The preferred route for medication administration should be based on the current patient condition and training level of the provider.					1 Glucagon calculated IV dose only in the event of Beta Blocker or Calcium Channel Blocker overdose.

Key:	
PO	Oral
SL	Sublingual
IN	Intranasal
IM	Intramuscular
IV	Intravenous
IO	Intraosseous





















